

# The Effectiveness of an Acceptance and Commitment Therapy Self-help Intervention for Chronic Pain

Marnie Johnston, MSocSci/PGDip Clin Psych,\* Mary Foster, PhD,\*  
 Jeannette Shennan, MSocSci/PGDip Clin Psych,† Nicola J. Starkey, PhD,\*  
 and Anders Johnson, MD, PhD†

**Objective:** To evaluate the effectiveness of an Acceptance Commitment Therapy based self-help book for people with chronic pain.

**Method:** This was a randomized 2 group study design. Over a 6-week period, 6 participants read the self-help book and completed exercises from it with weekly telephone support whereas 8 others formed a wait-list control group. Subsequently, 5 of the wait-list participants completed the intervention. Participants completed preintervention and postintervention questionnaires for acceptance, values illness, quality of life, satisfaction with life, depression, anxiety, and pain. Initial outcome data were collected for 8 control participants and 6 intervention participants. Including the wait-list controls, a total of 11 participants completed preintervention and postintervention measures. Whilst completing the self-help intervention, each week participants' rated the content of the book according to reading level and usefulness, and their comprehension of the content was also assessed.

**Results:** Compared with controls, participants who completed the book showed improved quality of life and decreased anxiety. When data from all the treatment participants were pooled, those who completed the intervention showed statistically significant improvements (with large effect sizes) for acceptance, quality of life, satisfaction with life, and values illness. Medium effect sizes were found for improvements in pain ratings.

**Conclusions:** These findings support the hypothesis that using the self-help book, with minimal therapist contact adds value to the lives of people who experience chronic pain.

**Key Words:** acceptance and commitment therapy, ACT, acceptance, chronic pain, self-help, quality of life, values.

(*Clin J Pain* 2010;26:393–402)

Chronic pain is problematic for those who experience it and can be complicated and costly to treat.<sup>1</sup> It often leads to disability and poor quality of life and is a costly social phenomenon. Psychological functioning and its relation to experiences of pain is a relatively recent area of inquiry. Before the development of the Gate Control Theory of Pain,<sup>2</sup> chronic pain was treated from a medical perspective with symptom reduction being the main aim.<sup>3</sup>

Subsequent developments in the field of pain management have resulted in biopsychosocial models of pain. Thus, current treatment approaches for chronic pain cover physical, psychological, and environmental factors and involve a multidisciplinary pain management approach that, although effective is expensive.

People with chronic pain often develop additional problems over time, such as emotional disturbance, cognitive difficulties, fear, and fatigue.<sup>4</sup> In such cases, cognitive-behavior therapy are often effectively used to change the client's focus from the pain itself, to the impact the pain is having on their functioning.<sup>5,6</sup> One recently developed Cognitive Behaviour Therapy is Acceptance and Commitment Therapy (ACT). The aim of ACT is to assist people to develop acceptance ("willingness to experience pain or distressing events without attempt to control them"<sup>7</sup>) of their pain and/or thoughts about their pain, together with enhancing the meaningful aspects of their lives (valued actions). Thus, rather than a futile struggle against pain, acceptance, and valued actions are used to improve quality of life, regardless of pain. There is growing evidence to support its effectiveness in the treatment of chronic pain.<sup>8</sup> However, the delivery of intensive individual therapeutic approaches, such as ACT, takes time, requires a skilled practitioner and although it is cheaper than treatment by multidisciplinary team, it can be costly. Therefore, the challenge is how to deliver such treatment programs effectively, tailored to individual needs, although minimizing time and cost.<sup>4</sup>

Self-help interventions are an approach that provide a means to administer standardized treatment to clients with minimal input from a therapist across a variety of disorders (eg, depression, anxiety disorders, eating disorders).<sup>9,10</sup> This approach has the additional benefits of being accessible to a larger numbers of clients at a much lower cost compared with other types of treatment. Thus, the overall aim of this study was to evaluate the effectiveness of an ACT-based self-help workbook "Living Beyond your Pain,"<sup>11</sup> (in combination with weekly phone and workbook support) for adults who experience chronic pain. More specifically, the study assessed whether participants would report "added value" to their lives after the intervention. "Added value" was considered an increase in acceptance, improvement in living in accordance with ones values and improvement in quality of life, depression, and anxiety.

## MATERIALS AND METHODS

### Settings and Participants

The study received approval from the Department of Psychology, University of Waikato, Ethics Committee, and

Received for publication September 17, 2009; revised October 22, 2009; accepted November 29, 2009.

From the \*Department of Psychology, Faculty of Arts and Social Sciences, The University of Waikato; and †Waikato Hospital Pain Clinic, Waikato District Health Board, Private Bag, Hamilton, New Zealand.

Reprints: Nicola J. Starkey, PhD, Department of Psychology, Faculty of Arts and Social Sciences, The University of Waikato, Private Bag, Hamilton, New Zealand (e-mail: nstarkey@waikato.ac.nz).

Copyright © 2010 by Lippincott Williams & Wilkins

the Northern Y District Health Board Ethics Committee. Potential participants were identified from the Waikato Hospital Pain Clinic Psychologist’s waiting list or were people who had been assessed by the Pain Clinic, but not referred to the psychologist, who as judged by the referral had a level of distress which might be helped by the intervention.

The inclusion criteria were; (1) Reading ability sufficient to read and understand the book, (2) No major psychiatric disorder that might impede participation, (3) Stable medication (eg, for 4 weeks and not considered by the Pain Clinic Psychologist to interfere with participation), (4) No known significant childhood history of trauma. These inclusion criteria were selected to ensure that participants would be able to complete the weekly reading and exercises, and to ensure their safety. Six people were excluded from the study owing to access difficulties and 2 others for their level of reading or cognitive ability.

After this, the Psychologist phoned 30 potential participants to determine if they were interested in participating in the study. Those who expressed an interest in the study were sent an information sheet and were contacted by the primary researcher.

Twenty-four participants volunteered thereafter to participate in the study and the majority of the meetings with these participants took place at their homes. Two participants were seen in the outpatient Pain Clinic rooms at Waikato Hospital. Of these, 37.5% (n = 10) of the participants were male and 62.5% (n = 14) were female. Ages ranged from 20 to 84 years with the median age being 43. The first participant was randomly designated to the control (C) or treatment (T) group and from then participants were assigned to each group alternately as they were contacted, to ensure that a similar number of participants were in each group. This resulted in 12 participants in each group.

Figure 1 illustrates the number of participants in each group and the attrition. Intention-to-treat analysis was applied for participants who withdrew after completing 3 weeks of the intervention (this was half of the treatment intervention and was considered sufficient for including in the posttest analysis). As can be seen in Figure 1, 5 of the initial treatment group completed the whole intervention; 1 participant withdrew after the third week of treatment, giving a subgroup of 6 who completed 3 or more weeks of treatment. Eight of the control group participants




FIGURE 1. Flow diagram of the study design and the fate of the participants in each group. Light gray boxes=control, dark gray=treatment.

completed the control period and provided predata and postdata. All of these participants agreed to start the treatment and are referred to as the C-T group. In this group, 5 completed the treatment. Thus, in total, 11 participated in at least 3 weeks of treatment.

## Measures

Table 1 shows outcome measures used in this study including number of items, and classification of scores for functionality. Corresponding domains of functioning as recommended by The Initiative on Methods, Measurement, and Pain Assessment in Clinical Trials<sup>12</sup> are also presented in Table 1.

Acceptance was measured by the Chronic Pain Acceptance Questionnaire (CPAQ). The CPAQ is a self-report questionnaire designed to measure acceptance of pain. It consists of 2 scales: activity engagement and pain willingness. Combined, the scales assess the individual's level of activity in the presence of pain and the degree to which the individual attempts to avoid or control pain. An increase in CPAQ score is associated with reported improvement in acceptance.<sup>13</sup>

Quality of life was used to evaluate participants' perceived subjective wellbeing. The Quality of Life Inventory (QOLI)<sup>14</sup> covers 17 domains of life that are evaluated by a weighted satisfaction score that includes satisfaction with the area of life and the importance of that area to the individual's well being. High scores on the QOLI correspond with a high level of perceived quality of life.

The Satisfaction with Life Scale (SWLS) is a brief self-report questionnaire that was designed to assess individual satisfaction with life as a whole.<sup>15</sup> This scale measures

the judgmental component (rather than the emotional component) of subjective well being; an increase in SWLS corresponds with improvement in an individual's judgment and reporting of their subjective wellbeing.

The Chronic Pain Values Inventory (CPVI) is a self-report questionnaire that was used to measure the concept *values illness* that is included in the self-help book. The first half of the items relate to specific areas of life and a rating is given by the individual to quantify the importance of each area. The latter half of the questions relate to how successful the individuals consider they have been at living in accordance with their values in each area. A discrepancy between importance and success indicates "values illness" and hence a decrease in CPVI score is associated with positive change for the individual.<sup>16</sup>

The Short-Form McGill Pain Questionnaire (SF-MPQ) is a self-report questionnaire that has been used to assess chronic pain experiences across a number of pain conditions. The questionnaire consists of 15 adjectives that describe sensory and affective aspects of pain in addition to a visual analogue scale and a present pain intensity (PPI) rating. High scores correspond with reports of high levels of pain.<sup>17</sup>

The Chicago Multi-scale Depression Inventory (CMDI) is a self-report inventory that was designed to measure depressive symptoms in medical patient populations.<sup>18</sup> The CMDI consists of 3 subscales; mood, evaluative, and vegetative which can be used individually or combined. The nonvegetative subscales are most accurate in examining depression in medical patients. High scores on the CMDI reflect reports of a high level of depression and decreases in CMDI score correspond with the reporting of a reduction in depressive symptoms.

**TABLE 1.** IMMPACT Criteria, the Corresponding Dependant Variables, and Measures Used in this Study With Classification Scores for Functionality

IMMPACT Criterion	Dependant Variable	Measure Name	Acronym	Number of Items	Classification of Scores
Physical functioning	Acceptance*	Chronic Pain Acceptance Questionnaire	CPAQ	20	Not applicable
Global improvement	Quality of life	Quality of Life Inventory	QOLI	32	< 42† 0-36 = very low, 37- 42 = low 43-57 = average, 58-7 = high
Global improvement	Satisfaction with life	Satisfaction with Life Scale	SWLS	5	Not applicable
Global improvement	Values illness*	Chronic Pain Values Inventory	CPVI	12	Not applicable
Pain	Pain	Short-Form McGill Pain Questionnaire	SF-MPQ	17	Present pain intensity only used 0 = no pain, 1 = mild, 2 = discomforting 3 = distressing 4 = horrible, 5 = excruciating
Emotional functioning	Depression	Chicago Multiscale Depression Inventory	CMDI	50	> 98†
Emotional functioning	Anxiety	Beck Anxiety Inventory	BAI	21	0-7 = min, 8-15 = mild 16-25 = moderate, 26-63 = severe

\*Not a formal/standardized measure.

†Manual provided cutoff score for functionality.

**TABLE 2.** Details of the Book Chapters Covered Each Week and a Summary of the Content

Week	Chapters	Summary of Content
1	Introduction Ch 1 & 2	<i>What is pain? What is ACT?</i> Changing one's relationship with pain, the main ideas of ACT, the pain avoidance suffering cycle, how one's experience of pain may limit participation in valued activities.
2	Ch 3	<i>What do you value?</i> Identifying values (important aspects of life) and the concept that values are a way of living. Introduction of 'values illness'-how one's experience with pain may inhibit them from living a life they value
3	Ch 4	<i>You are not your thoughts</i> How to develop distance from thoughts. Introduces cognitive defusion and the idea of thought observation without judgment.
4	Ch 5	<i>Mindfulness</i> Identifying 3 aspects of the self, and who are we if our thoughts don't identify us? How to practice mindfulness.
5	Ch 6, half Ch 7	<i>Are you willing? Committed actions</i> Introduces acceptance and willingness to engage with pain. Taking committed actions (deciding what to do and doing it), and identifying obstacles to this.
6	Remainder Ch 7, Ch 8	<i>Committed actions &amp; what's holding you back?</i> Identifying vital (actions that support living a valued life) and nonvital actions, using the bulls-eye to get closer to values. Thinking about future obstacles and developing committed action plans for obstacles which may be encountered in the future

The Beck Anxiety Inventory (BAI) is a self-report questionnaire that was designed to measure symptoms of anxiety.<sup>19</sup> This measure was used in addition to the CMDI as a measure of emotional functioning. High scores on the BAI reflect high level of anxiety symptoms reported by the individual and a reduction in score corresponds with reported improvement in anxiety symptoms. Half of the items on the BAI represent somatic symptoms of anxiety and the remainder represents cognitive symptoms.

## Materials

Each participant who undertook treatment was provided with a copy of Dahl and Lundgren (2006)<sup>11</sup> book, "Living Beyond Your Pain," together with a workbook. The book is composed of 8 chapters, in addition to introduction and conclusion sections. The chapters and topics covered in each week are summarized in Table 2. The workbook was compiled by the primary researcher and was developed to be used alongside the text. It included a front sheet explaining the weekly intervention requirements that were to read the specified part of the book, complete the exercises, and answer questions about these during the weekly phone call. Participants were reassured that the questions during the phone call were not a test. The remainder of the workbook was divided into 6 sections, with 1 to be completed per week. Each section consisted of a summary of the week's reading, spaces to write and complete exercises from Dahl and Lundgren (2006), and a list of question relating to the content for that week. The full contents of the workbook can be viewed at <http://hdl.handle.net/10289/2309>.

## Procedure

Once potential participants had received the information sheet in the mail, the primary researcher phoned them to discuss the study details and their participation. For those who wanted to take part, an initial meeting was arranged at either the Pain Clinic or at the participant's home. As described above, the first participant was randomly desig-

nated to the control or treatment group and from then participants were assigned to each group alternately as they were contacted. For all participants, the first meeting consisted of reviewing the information sheet, explaining the study, and signing the consent form. The battery of tests was completed and a weekly time was arranged for the researcher to make phone contact with the participant.

## Control Group

At the initial meeting, in addition to completing the tests and arranging a telephone interview time, the control group participants were offered a second meeting 6 weeks later to repeat the battery of tests and have the opportunity to start the intervention. Control participants were phoned by the primary researcher once a week on the prearranged day and time. During each phone call, they were asked to rate their pain on a 10-point scale in which 0 is no pain and 10 is the worst pain imaginable. They also rated their pain according to the Present Pain Intensity (PPI) from the SF-MPQ and were asked if there was anything that they perceived was influencing their pain in a positive or negative way. The PPI is a 6-item scale in which 0 is no pain and 5 is excruciating pain. The numbers between 0 and 5 have corresponding words to describe the level of pain. The main purpose of these questions for the Control group was to establish similar conditions for the Treatment and Control groups, that is, participants from both groups received weekly phone calls and discussed issues around pain. The weekly data from the Control group were not analyzed further.

After 6 weeks of weekly contact, participants repeated the battery of tests and were offered the opportunity to continue to the intervention. Participants that moved to the intervention phase followed the procedure described below. After completing the intervention, these participants completed the battery of tests for the third time.

## Treatment Group

Treatment participants were loaned a copy of the self-help book and given their own copy of the accompanying

workbook to use during the intervention and keep afterward. Participants were instructed to start the first week of the intervention right away. This involved them reading the selected portion of the text (documented in the front sheet of the workbook), completing the exercises in the workbook for that section, and noting the answers to the questions for that week (Table 2).

The weekly phone calls from the primary researcher to these participants involved 3 standard questions: 1. Did you do all, some, or none of the reading and the exercises? 2. Did you find the reading level easy, medium, or hard? 3. Was the book very useful, moderately useful, or not useful at all? Once these questions had been responded to, the researcher addressed the questions in the workbook for each week. The participants relayed their answers to the researcher who wrote down the participants' responses verbatim. These answers were later evaluated by the researcher to assess comprehension of the material described in the book.

After they had completed reading the book and using the workbook over a 6-week period, Treatment participants repeated the battery of tests and thereby completed the study. The dependant variables measured by the battery of tests included acceptance, subjective wellbeing, values illness, pain, and emotional functioning.

## RESULTS

### Statistical Analysis

Data Analyses were conducted using the Statistical Package for the Social Sciences (SPSS) version 14.0. Once data collection was completed, *t* tests were used to compare participants assigned with the control and treatment groups on all measures. Thereafter, statistical analyses were

conducted using *t* tests and/or ANOVA's as appropriate. The  $\alpha$  level for statistical significance was defined as  $P < 0.05$ .

As the sample size was small, statistical effect sizes were included to give a measure of the magnitude of change (from preintervention to postintervention) on each measure. Effect size provides a way to analyze the importance of change irrespective of sample size. Statistical effect sizes were calculated as part of the ANOVA (partial  $\eta^2$ ) and were considered small if between 0.10 to 0.29; medium if between 0.30 and 0.49; and large if greater than 0.50.<sup>20</sup>

As well as looking at looking at statistical effects, clinically significant improvement was examined. Clinical significance is defined as a change in reported level of functioning that corresponds with a shift from a clinical population (before treatment) to a general (more functional) population (after treatment).<sup>21</sup> A shift in score toward the mean of the general population is considered to show improvement. The range of measures used in this study meant that clinically significant improvement had to be defined in different ways (Table 1). For measures (such as the ACT measures) in which no clinical cutoff was reported in the literature, improvement in score after treatment, was considered clinically significant improvement. The cutoffs for the QOLI and CMDI were defined by using their respective manual specifications for symptoms in the clinical range. For pain, PPI scale from the SF-MPQ was used to analyze clinically significant change and is shown as any movement in the direction of improvement.

### Initial Analysis: Control Versus Treatment

The means and standard deviations of the scores for the tests from the first administration (T1) for all of the initial treatment (n = 6) and control groups (n = 8) and from the first and second administrations (T2) for those

**TABLE 3.** Means, Standard Deviations (SD), and Number of Participants (N) for Each Test in the Battery for all Participants at Time 1, and for Those Who Completed to Time 2, at Time 1, and Time 2

Group	Time 1 (T1)						Time 2 (T2)		
	All Participants Data at T1			T1 Data for Participants Who Completed to T2			T2 Data for Participants Who Completed to T2		
	N	Mean	SD	N	Mean	SD	N	Mean	SD
Chronic Pain Acceptance Questionnaire (CPAQ)									
Control	11	56.3	17.0	8	52.3	13.8	8	62.3	24.8
Treatment	12	49.5	13.1	6	59.0	8.2	5	77.4	9.1
Quality of Life Inventory (QOLI)									
Control	11	35.3	16.7	8	31.1	12.6	8	34.4	19.6
Treatment	12	35.8	11.7	6	41.5	9.7	6	51.0	5.6
Satisfaction with Life Scale (SWLS)									
Control	11	17.6	6.9	8	17.4	4.0	8	18.1	8.3
Treatment	12	15.7	6.0	6	19.3	6.5	6	22.5	6.5
Chronic Pain Values Inventory (CPVI)									
Control	11	10.0	7.7	8	16.0	9.4	8	7.0	8.7
Treatment	12	11.3	7.7	6	12.7	4.8	6	3.5	3.0
McGill Pain Questionnaire, Short Form (SF-MPQ)									
Control	11	20.5	11.9	8	22.4	10.4	8	20.1	8.4
Treatment	12	21.2	10.9	6	16.0	9.6	5	15.6	7.5
Chicago Multiscale Depression Inventory (CMDI)									
Control	11	104.4	31.0	8	113.9	24.4	8	102.6	25.0
Treatment	12	122.6	40.9	6	89.0	23.9	6	86.3	25.6
Beck Anxiety Inventory-II (BAI)									
Control	11	15.8	8.9	8	20.0	8.8	8	18.5	7.7
Treatment	12	17.5	13.5	6	12.0	8.7	6	8.3	5.4

**TABLE 4.** Two-way ANOVA and Independent *t* test Results Evaluating the Effects of the Self Help Intervention on the Initial Sample

Measure	df	ANOVA						<i>t</i> -tests (Control vs. Treatment)					
		Group (G) (Treatment/Control)		Time (T1/T2)		Interaction (G×Time)		T1			T2		
		<i>F</i>	<i>P</i> η <sup>2</sup>	<i>F</i>	<i>P</i> η <sup>2</sup>	<i>F</i>	<i>P</i> η <sup>2</sup>	<i>t</i>	<i>df</i>	<i>d</i>	<i>t</i>	<i>df</i>	<i>d</i>
CPAQ	1, 11	1.62	0.13	9.61*	0.47	0.90	0.08	-1.06	12	<b>0.61</b>	-1.57	9.5	<b>1.0</b>
QOLI	1, 12	4.33	0.27	3.55	0.23	0.85	0.07	-1.68	12	<b>0.97</b>	-2.29*	8.4	<b>1.6</b>
SWLS	1, 12	1.00	0.08	1.66	0.12	0.63	0.05	-0.70	12	<i>0.40</i>	0.38	12	0.22
CPVI	1, 12	1.09	0.08	15.74*	0.57	0.01	0.01	0.79	12	<i>0.46</i>	0.93	12	<i>0.54</i>
SF-MPQ	1, 11	0.85	0.07	1.28	0.10	0.03	0.01	1.16	12	<b>0.67</b>	0.98	11	<i>0.59</i>
CMDI	1, 12	2.78	0.19	1.92	0.14	0.73	0.06	1.91	12	<b>1.10</b>	1.20	12	<i>0.69</i>
BAI	1, 12	9.14*	0.43	0.75	0.06	0.13	0.01	1.69	12	<b>0.98</b>	2.76*	12	<b>1.6</b>

\**P* < 0.05.

BAI indicates Beck Anxiety Inventory; CMDI, Chicago Multiscale Depression Inventory; CPAQ, Chronic Pain Acceptance Questionnaire; CPVI, Chronic Pain Values Inventory; *d*, Cohen's effect size (**bold** = large, *italics* = medium); *df*, degrees of freedom; *F*, ANOVA statistic; MPQ-SF, Short-Form McGill Pain Questionnaire; *P* η<sup>2</sup>, partial eta squared; QOLI, Quality of Life Inventory; SWLS, Satisfaction With Life Scale; *t*, *t* obtained from independent groups *t*-test.

who completed both the premeasures and postmeasures are given in Table 3. The original Control and Treatment groups (all participants) obtained similar scores on most measures apart from the CMDI. However, when looking at the baseline measures for participants who completed to T2, differences between the scores of the control and treatment groups at baseline are evident for most measures. Six weeks later (T2), both the control and treatment groups have shown improvements on all measures.

Two-way repeated measures ANOVAs (groups: control and treatment; time: T1 and T2) were conducted to compare data for those who completed both T1 (pre) and T2 (post) measures. The results of these are given in the first 3 columns of Table 4. In Table 4, and thereafter, statistically significant values are asterisked, bolded values represent large effect sizes, and italicized values represent medium effect size using the criteria suggested by Cohen.<sup>20</sup>

From Table 4 it can be seen that the main effect of group membership was significant for the BAI, with the controls showing higher levels of anxiety compared with the treatment group. The main effect of time of administration (T1/T2) was significant for the CPAQ and the CPVI, reflecting the increase in acceptance scores and decrease in values illness scores over the 6 week period in both groups. All other effects were not statistically significant.

Table 4 also summarizes the results of independent *t* tests comparing the control and treatment groups test

scores at T1 and T2. These show the T1 scores were not significantly different between the groups. However, at T2 there were significant differences between the groups for scores on the QOLI and BAI (with large effect sizes). The treatment group showed significantly improved quality of life and lower levels of anxiety compared with controls.

The scores from the other 5 tests CPAQ, SWLS, CMDI, CPVI, and MPQ-SF were not significantly different between the groups at T2. However, large effect sizes were found for the difference in CPAQ scores (treatment group showing higher acceptance), and medium effect sizes for CPVI, CMDI, and MPQ-SF (the treatment group showing more favorable scores in each case). The effect size for SWLS was small.

**Second Level of Analysis: Control-Treatment (C-T) Group**

The means and SEs of the scores from all test administrations for each test for the 5 control participants who continued with the treatment (and completed a minimum of 3 weeks of the intervention) and are given in Table 5. The results of repeated measures *t* tests, comparing the measures between T1 and T2 and between T2 and T3 for the C-T group, are also in Table 5. These data show that whereas there were no significant differences between the T1 and T2 or T2 and T3 administrations, there were large effect sizes for T2 versus T3 for the QOLI, SWLS, and

**TABLE 5.** The Mean Scores Obtained by the Control-Treatment Group Participants, at the Start of the Control Period (T1), the Beginning of the Intervention (T2), and After Completing the Intervention (T3)

Measure	N	Mean (SE)			T1 vs. T2			T2 vs. T3		
		T1	T2	T3	<i>t</i>	<i>df</i>	<i>d</i>	<i>t</i>	<i>df</i>	<i>d</i>
CPAQ	5	52.6 (9.3)	54.0 (7.9)	72 (11.6)	-0.5	4	0.22	-2.7	4	0.01
QOLI	5	33.8 (10.3)	33.8 (6.9)	43.8 (6.7)	0.1	4	0	-2.0	4	<b>0.89</b>
SWLS	5	16.0 (4.2)	18.2 (2.2)	22.4 (3.0)	-0.8	4	0.34	-2.5	4	<b>1.1</b>
CPVI	5	11.0 (2.9)	16.0 (5.2)	4.4 (3.4)	-0.8	4	0.37	2.2	4	<b>0.96</b>
MPQ-SF	5	22.0 (3.6)	22.8 (2.1)	19.8 (3.0)	-0.4	4	0.17	1.6	4	<i>0.73</i>
CMDI	5	113.1 (10.9)	111.7 (4.6)	107.1 (12.1)	0.2	4	0.08	0.5	4	0.22
BAI	5	20.0 (7.0)	21.8 (3.2)	19 (4.2)	-0.4	4	0.19	0.4	4	0.17

Data are presented as mean (SE), the *t* values From dependent *t*-Tests comparing across the time periods are also presented.

BAI Indicates Beck Anxiety Inventory; CMDI, Chicago Multiscale Depression Inventory; CPAQ, Chronic Pain Acceptance Questionnaire; CPVI, Chronic Pain Values Inventory; *d*, Cohen's *d* effect size (**bold** = large, *italics* = medium); *df*, degrees of freedom; SF-MPQ, Short-Form McGill Pain Questionnaire; N, number of participants; QOLI, Quality of Life Inventory; SWLS, Satisfaction With Life Scale; T1, time 1; T2, time 2; T3, time 3.

**TABLE 6.** The Effect of the Self-help Book on Test Scores From Participants Who Completed at Least 3 Weeks of Treatment

Measure	N	Mean (SE)		Pretreatment vs. Posttreatment <i>t</i> Tests			Clinical Significance
		PreTest	PostTest	<i>t</i>	<i>df</i>	<i>d</i>	No. Participants With Clin. sig. Improvement
CPAQ	10	56.3 (4.3)	74.7 (5.9)	-4.1*	9	<b>1.30</b>	9
QOLI	11	38 (3.8)	47.7 (3.3)	-3.63*	10	<b>1.10</b>	10 (4 of which above cut-off)
SWLS	11	18.8 (1.7)	22.5 (1.9)	-2.82*	10	<b>0.85</b>	10
CPVI	11	14.2 (2.5)	3.9 (1.6)	4.04*	10	<b>1.21</b>	10
SF-MPQ	10	20.7 (2.1)	17.7 (2.2)	1.73	9	0.55	4
CMDI	11	99.3 (6.5)	95.8 (8.2)	0.71	10	0.21	6 (1 of which below cut-off)
BAI	11	16.5 (2.8)	13.2 (2.7)	0.95	10	0.29	7 (3 of which moved categories)

Data are presented as mean (SE) along with the results of the repeated measures *t* test and summary of clinically significant improvement.

\**P* < 0.05.

BAI indicates Beck Anxiety Inventory; CMDI, Chicago Multiscale Depression Inventory; CPAQ, Chronic Pain Acceptance Questionnaire; CPVI, Chronic Pain Values Inventory; *d*, Cohen's *d* effect size (**bold** = large, *italics* = medium); *df*, degrees of freedom; SF-MPQ, Short-Form McGill Pain Questionnaire; N, number of participants; QOLI, Quality of Life Inventory; SWLS, Satisfaction With Life Scale.

CPVI, and medium effect size for MPQ-SF. A small effect size was seen for the CMDI for T2 versus T3.

**Third Level of Analysis: Pooled Data From All Participants Completing the Intervention**

Table 6 gives the means and SEs for all the tests for participants who completed at least 3 weeks of the treatment. This includes data from the initial analysis together with the C-T group's T2 (preself-help intervention) and T3 (postself-help intervention) data. The pretest scores are those from T1 for the initial treatment group and from T2 for the C-T group. The posttests scores are those from T2 for the initial treatment group and from T3 for the C-T group. The results of paired *t* tests (pretreatment and posttreatment) for all treatment completers are given in Table 6, whereas the percentage change in each of the measures can be seen in Figure 2.

Statistically significant differences (and large effects sizes) were found between the preintervention and post-


intervention test scores for CPAQ, QOLI, SWLS, and CPVI. This shows that overall; there were improvements in acceptance, quality of life, satisfaction with life, and values illness for those who completed at least 3 weeks of the self-help intervention. This is reflected by the large percentage change in these scores which can be seen in Figure 2. Furthermore, a medium effect size was observed for the change in score on the SF-MPQ and a small effect size was found for the CMDI.

For clinical significance, Table 6 shows the number of participants for whom there were clinically significant improvements (as earlier defined) for each measure on completion of the self-help intervention. As can be seen in Table 6, the majority of participants' quality of life, satisfaction with life, acceptance, and values illness measures showed clinically significant improvements. Just over half of the participants' anxiety and depression measures showed improvement, and for pain 4 people reported decreased pain intensity on completion of the intervention.

**Ratings of the Self-help Book**

To ensure treatment integrity and to provide information about the way participants experienced the self-help book, data were collected during the weekly phone calls from participants who were completing the treatment. These data are shown in Table 7. Participants rated the degree to which they completed the weekly requirement (presented as percentage of participants who completed all, some or none of the weekly requirements). Comprehension scores (derived from weekly questions asked on the phone) were given to each participant per week and are presented in Table 7. This measure indicates the percentage of questions about the material for that week answered correctly during the weekly phone call.

Participants also rated how useful they found the weekly requirements (percentage of participants who rated the weekly material of no, some, or high use), and how difficult they found the reading level for the weekly reading (percentage of participants that rated the reading level easy, medium, or hard). These data are also included in Table 7. As can be seen in Table 7, most participants completed all or some of the readings and exercises for each week. In terms of comprehension, the contents of week 4 (mindfulness) seemed to be less well understood by the participants in this study. Participants' ratings for usefulness of the weekly



**FIGURE 2.** Percent change (+SE) from pretreatment to post-intervention for all participants completing more than 3 weeks of the self help book. See text for further details. BAI indicates Beck Anxiety Inventory; CMDI, Chicago Multiscale Depression Inventory; CPAQ, Chronic Pain Acceptance Questionnaire; CPVI, Chronic Pain Values Inventory; SF-MPQ, Short-Form McGill Pain Questionnaire; QOLI, Quality of Life Inventory; SWLS, Satisfaction With Life Scale.

**TABLE 7.** Participant's Weekly Ratings for Completion of Material, Utility of Material, Difficulty of Material, and Comprehension of Material, Obtained From Weekly Telephone Support

Week	1	2	3	4	5	6
Corresponding Content of the Book	Definitions of Pain & ACT Control is not the Answer	Values	Cognitive Defusion	Mindfulness	Willingness & Committed Action	Committed Action & Obstacles
Completion (%)						
All	81	79	75	56	78	67
Some	13	14	25	44	22	33
None	6	7	0	0	0	0
Comprehension (% score)						
Mean	72	79	73	58	77	81
Sd	21	30	19	22	16	13
Use (%)						
Very	50	39	50	36	56	33
Medium	43	46	25	45	22	44
None	7	15	25	18	22	22
Difficulty (%)						
Easy	33	43	25	22	33	56
Medium	53	36	33	44	44	22
Hard	13	21	41	33	22	22

reading and exercises show variation, with the highest number of “no use” ratings for week 3 (cognitive defusion). For difficulty ratings, week 3 (cognitive defusion) received the most “hard” ratings, and the highest number of “easy” ratings were given for weeks 2 (values) and 6 (committed action).

## DISCUSSION

The results of this study support the effectiveness of the self-help book by Dahl and Lundgren<sup>11</sup> for adults with chronic pain. In particular, statistically and clinically significant improvement and large effect sizes were shown for participants' acceptance, quality of life, satisfaction with life, and values illness scores. In addition, the pooled data indicated a medium effect size for participants' reports of reduced pain after they had completed the intervention. Thus, this study showed that adult participants with chronic pain benefited from using an ACT-based self help manual, with therapist support. These findings are in keeping with those reported in Watkins and Clum<sup>10</sup> review of self-help interventions and they provide support for earlier studies which showed the efficacy of self-help type interventions with minimal therapist contact, for children with chronic headache and recurrent pain.<sup>22,23</sup>

Acceptance is a core process of ACT and one-to-one ACT therapy increases acceptance.<sup>24-26</sup> Acceptance also forms a central theme of Dahl and Lundgren's book<sup>11</sup> and given the significant increases in acceptance reported here, this suggests that ACT can be effectively delivered through a self-help book.

From an ACT perspective acceptance is not passive but involves allowing one to experience thoughts, feelings, and physical sensations as a means to engage in valued activities.<sup>27</sup> In this study activity levels were not formally measured, however, many of the current participants reported increasing their involvement in activities they valued. These changes may be the result of the emphasis ACT gives to “willingness to engage” and the text's<sup>11</sup> aim to increase the reader's valued activities and to minimize their avoidance behavior.

Acceptance of pain, and engaging in activity despite pain, have been found to correlate positively with quality of life.<sup>28,29</sup> In this study those who participated in the self-help intervention reported higher quality of life and satisfaction with life at the end of the intervention. These findings add to those from acceptance and mindfulness-based studies with problems other than pain (eg, with Generalized Anxiety Disorder<sup>30</sup>) which have found increases in subjective well-being. The increases in subjective wellbeing reflect participants' reports of increased satisfaction with many aspects of their lives.

The role of values is also a core process of ACT and is emphasized by Dahl and Lundgren.<sup>11</sup> The efficacy of using one's personal values to assist with living with pain has been well-established.<sup>16,31,32</sup> In support of these findings, this study found significant improvement as a result of treatment in terms of living in alignment with values. McCracken and Vowles<sup>32</sup> propose that acceptance and values-based action are distinct but related processes and the improvement shown in both of these areas by participants in this study support this view.

From an ACT perspective, the focus of therapy is not on reducing pain but on living a valued life. These results showed that although participants increased their acceptance of living with pain, their pain (SF-MPQ) scores did not decrease significantly. Given that overall there was a medium statistical effect size for reduction of pain; the sample size may have influenced this outcome. These findings were similar to those of Morone, Greco, and Weiner<sup>33</sup> who carried out a randomized controlled pilot study to evaluate an 8-week mindfulness group intervention for community dwelling older adults with lower back pain. Using the same measure (SF-MPQ), they found that participants from their intervention group showed improvement in pain scores on completion of the intervention, but this did not reach statistical significance. Interestingly, the Morone study<sup>33</sup> also reported statistically significant improvement in acceptance scores (also measured by the CPAQ) for the mindfulness intervention group. Other studies have reported statistically significant reductions in pain severity when ACT<sup>7</sup> or components of ACT are

used.<sup>34,35</sup> However, these studies used different measures, different populations, and a different treatment format. Using the PPI rating scale as a way to assess clinically significant change in pain, showed that just under half the participant's reported some improvement in pain once they had completed the intervention. In this study, pain scores may have been influenced by participants' use of pain medication at data collection times (when the SF-MPQ was administered). It was common for participants to comment that they had taken pain medications a short time prior and that their pain had been worse at a different time of the day. Such factors are likely to have interfered with the accuracy of the predata and postdata collected for participants' pain.

Similar to other outcomes reported in the literature,<sup>36</sup> several participants from this study commented on the difficulty in engaging with the intervention when their pain level was higher than usual. In line with these reports, those who pulled out of the study reported higher initial pain score. However, those who participated still experienced pain.

There was little evidence to support the notion that ACT therapy delivered through a self help book improved mood. This is contrary to other research.<sup>26,37-40</sup> This may, in part, be owing to the use of different measures to assess mood. For example, the Pain Anxiety Symptom Scale and the BDI were most commonly used in the above studies, rather than the BAI and the CMDI, as used in this study. The BAI was chosen for this study to provide an overall measure of anxiety, rather than one which is pain specific, whilst the CMDI was used as it minimizes the overlap between symptoms of depression and symptoms associated with pain. At baseline, around half of the participants obtained scores within the normal or nonclinical range for anxiety and depression. Thus, this group would not necessarily be expected to improve further. For example, on the CMDI, 5 of the 11 participants who completed the intervention scored below 98 (the cut off score) at baseline. Anxiety scores also showed this pattern. However, the majority of participants whose scores reflected severe depression and anxiety preintervention had improved (at least one level, eg, moderate to mild for anxiety or below the cut-off score for depression) at postintervention. That is even though the changes in these measures were not statistically significant, overall, the treatment completers who scored in the "dysfunctional" range before treatment showed clinically significant improvements in depression and anxiety scores on completion of treatment. As cautioned by Jacobson and Truax<sup>21</sup> cutoff scores can be arbitrary and may not capture all improvements (eg, from one end of a range to another). One participant's anxiety score was 14 (high end of mild range) at baseline and 8 postintervention (cut off from minimum to mild range), illustrating the way in which even an analysis of clinical significance may not reflect the full picture of improvement. With this in mind, we conclude that the majority of participants who reported problematic anxiety at baseline showed clinically significant improvement after completing the intervention.

Cuijpers<sup>9</sup> emphasized the importance of the role of the professional in supported self-help interventions and this study led to questions about how to define therapist "support" participants received from the researcher. For example, the role of the researcher involved more than data collection but less than regular one-on-one therapy. Throughout the intervention, participants commented on the importance of the weekly phone call for motivation

to complete the weekly tasks and in giving them the opportunity to ask questions. On a small number of occasions therapist input was required to support participants with difficulties that arose during the process of the intervention. This outcome mirrors the existing literature that documents the benefits of providing therapist support with a self-help intervention.<sup>9,41</sup> These outcomes also support Watkins and Clums'<sup>10</sup> proposition that the role of the practitioner be made explicit to intervention participants.

### Limitations

This study used a small sample and included a high proportion of participants who either did not start or withdrew from the intervention. The numbers of those originally contacted to ascertain interest in participating in the study were not officially recorded. Thus the flow chart of participants begins with the 24 people who agreed to participate. However, the fact that the treatment produced statistically significant effects in this sample adds more weight to the current findings, as it is more difficult to obtain statistically significant findings with a small number of participants.

In addition, no longer term follow-up data were obtained to evaluate the persistence of the effects. Owing to variation in the measures used in the study, a consistent analysis of clinically significant change was difficult. To provide consistency in discussing clinically significant change, and as recommended by Jacobson and Truax,<sup>21</sup> overall improvement (shift toward mean of general population) was reported. In addition to this, information about movement from dysfunctional classification of score to more functional classification of score was also included when possible. Additional measures, for example, of readiness to change and of level of disability, could have been included in this study; however, adherence to the IMMPACT recommendations was considered sufficient.

### Summary

This study shows support for the use of a self-help intervention with therapist support, for people with chronic pain. Those who completed the intervention showed improved acceptance, quality of life, and satisfaction with life in addition to living more closely aligned with their values. Further research is warranted with a larger sample size to assess the longer term effects of the intervention, as it may prove to be a cost-effective alternative for treatment of chronic pain. Such research could include an analysis of the characteristics of those who might benefit from such an approach. In addition, it would be useful to examine the role of the therapist in the delivery and outcomes for self-help participants.

### ACKNOWLEDGMENT

*The authors thank the Waikato Hospital Pain Clinic, Hamilton, New Zealand, and those who participated in the study.*

### REFERENCES

1. Gatchel RJ, Peng YB, Peters ML, et al. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychological Bulletin*. 2007;133:581-624.
2. Melzack R, Wall PD. *The Challenge of Pain: a Modern Medical Classic*. London, United Kingdom: Penguin Books; 1996.
3. Fordyce WE. Pain and suffering: a reappraisal. *Amer Psychologist*. 1988;43:276-283.

4. Eccleston C. Role of psychology in pain management. *Brit J Anaesth*. 2001;87:144–52.
5. Lohnberg JA. A review of outcome studies on cognitive-behavior therapy for reducing fear-avoidance beliefs among individuals with chronic pain. *J Clin Psychological Med Settings*. 2007;14:113–122.
6. Morley S, Eccleston C, Williams A. Systematic review and meta-analysis of randomized controlled trials of cognitive behaviour therapy for chronic pain in adults, excluding headache. *Pain*. 1999;80:1–13.
7. Vowles KE, McCracken LM. Acceptance and values-based action in chronic pain: a study of treatment effectiveness and process. *J Consul Clin Psychol*. 2008;76:397–407.
8. Dahl J, Wilson KG, Luciano C. *Acceptance and Commitment Therapy for Chronic Pain*. Reno, NV: Context Press; 2005.
9. Cuijpers P. Bibliotherapy in unipolar depression: a meta-analysis. *J Beha Ther Expl Psychiatry*. 1997;28:139–147.
10. Watkins PL, Clum GA, (eds). *Handbook of Self Help Therapies*. New York, US: Routledge/Taylor & Francis Group; 2008.
11. Dahl J, Lundgren T. *Living Beyond Your Pain: Using Acceptance and Commitment Therapy to Ease Chronic Pain*. Oakland, America: New Harbinger Publications, Inc; 2006.
12. Dworkin RH, Turk DC, Farrar JT, et al. Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain*. 2005;113:9–19.
13. McCracken LM, Vowles KE, Eccleston C. Acceptance of chronic pain: component analysis and a revised assessment method. *Pain*. 2004;107:159–166.
14. Frisch MB, Cornell J, Villanueva M, et al. Clinical evaluation of the quality of life inventory: a measure of life satisfaction for use in treatment planning and outcome assessment. *Psychological Assessment*. 1992;4:92–101.
15. Weinman J, Wright S, Johnston M. *Health Status and Health-Related Quality of Life*. Windsor, UK: Nfer-Nelson; 1995.
16. McCracken LM, Yang S. The role of values in a contextual cognitive-behavioral approach to chronic pain. *Pain*. 2006;123:137–145.
17. Melzack R. The short-form McGill pain questionnaire. *Pain*. 1987;30:191–197.
18. Nyenhuis DL, Luchetta T, Yamamoto C. The development, standardisation, and initial validation of the Chicago multi-scale depression inventory. *J Personality Assessment*. 1998;70:386–401.
19. Beck AT, Epstein N, Brown G, et al. An inventory for measuring clinical anxiety: psychometric properties. *J Consulting Clin Psychol*. 1988;56:893–897.
20. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2th ed. New Jersey: Lawrence Erlbaum Associates Inc; 1988.
21. Jacobson NS, Truax P. Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *J Consulting Clin Psychol*. 1991;39:12–19.
22. Griffiths JD, Martin PR. Clinical- versus home-based treatment formats for children with chronic headache. *Brit J Health Psychol* 1996;1:151–166.
23. Hicks CL, von Baeyer CL, McGrath PJ. Online psychological treatment for pediatric recurrent pain: a randomized evaluation. *J Pediatric Psychol*. 2006;31:724–736.
24. Forman EM, Herbert JD, Moitra E, et al. A randomized controlled effectiveness trial of acceptance and commitment therapy and cognitive therapy for anxiety and depression. *Behavior Modification*. 2007;31:772–799.
25. McCracken LM, Eccleston C. Coping or acceptance: what to do about chronic pain? *Pain*. 2003;105:197–204.
26. McCracken LM, MacKichan F, Eccleston C. Contextual cognitive-behavioral therapy for severely disabled chronic pain sufferers: effectiveness and clinically significant change. *Euro J Pain*. 2007;11:314–322.
27. McCracken LM, Carson JW, Eccleston C, Keefe FJ. Acceptance and change in the context of chronic pain. *Pain*. 2004;109:4–7.
28. Kratz AL, Davis MC, Zautra AJ. Pain acceptance moderates the relation between pain and negative affect in female osteoarthritis and fibromyalgia patients. *Behavioral Med*. 2007;33:291–301.
29. Mason VL, Mathias B, Skevington SM. Accepting low back pain: is it related to a good quality of life? *Clin J Pain*. 2008;24:22–29.
30. Roemer L, Orsillo SM. An open trial of an acceptance-based behavior therapy for generalized anxiety disorder. *Behavior Therapy*. 2007;38:72–85.
31. McCracken LM. *Contextual Cognitive-Behavioral Therapy for Chronic Pain: Progress in Pain Research and Management, volume 33*. Seattle: IASP Press; 2005.
32. McCracken LM, Vowles KE. A prospective analysis of acceptance of pain and values-based action in patients with chronic pain. *Health Psychol*. 2008;27:215–220.
33. Morone NE, Greco CM, Weiner DK. Mindfulness meditation for the treatment of chronic low back pain in older adults: a randomized controlled pilot study. *Pain*. 2008;134:310–319.
34. Baer RA. Mindfulness training as a clinical intervention: a conceptual and empirical review. *Clin Psychol Science Practice*. 2003;10:125–143.
35. Kabat-Zinn J, Lipworth L, Burney R. The clinical use of mindfulness meditation for the self regulation of chronic pain. *J Behavioral Med*. 1985;8:163–190.
36. Jensen M, Turner J, Romano J, et al. Coping with chronic pain: a critical review of the literature. *Pain*. 1991;47:249–283.
37. Dalrymple KL, Herbert JD. Acceptance and commitment therapy for generalized social anxiety disorder: a pilot study. *Behaviour Modification*. 2007;31:543–568.
38. McCracken LM, Gauntlett-Gilbert J, Vowles K. The role of mindfulness in a contextual cognitive-behavioural analysis of chronic pain-related suffering and disability. *Pain*. 2007;131:63–69.
39. McCracken LM. Learning to live with pain: acceptance of pain predicts adjustment in persons with chronic pain. *Pain*. 1998;74:21–27.
40. Roemer L, Orsillo SM, Salters-Pedneault K. Efficacy of an acceptance-based behavior therapy for generalized anxiety disorder: evaluation in a randomised controlled trial. *J Consulting Clin Psychol*. 2008;76:1083–1089.
41. Floyd M, Rohen N, Shackelford JAM, et al. Two-year follow-up of bibliotherapy and individual cognitive therapy for depressed older adults. *Behavior Modification*. 2006;30:281–294.